

UBC Physics Circle



Session 2: Problems


November 22, 2023

For our second problem session of 2023, we will follow in the illustrious footsteps of Stephen Hawking, and discover that black holes *glow*. This means they slowly leak energy into space, and eventually vanish in a burst of high-energy radiation! We will use this to calculate the lifetime of a solar mass black hole.


We're going to need five basic facts from Douglas Scott's talk. These are:

- the size of a black hole;
- the uncertainty principle of quantum mechanics;
- the relativistic energy of a moving particle;
- the average energy of particles in hot systems; and
- the rate a hot object radiates energy.


We picture these using cartoons below.




$$R_s = \frac{2GM}{c^2}$$




$$\Delta x \Delta p \geq \frac{h}{4\pi}$$



$$E^2 = p^2 c^2 + m^2 c^4$$



$$E_{\text{avg}} \sim k_B T$$



$$L = \sigma A T^4$$

Let's go through these more carefully:

- **Black holes.** First, a black hole of mass M has a *Schwarzschild radius* of

$$R_s = \frac{2GM}{c^2}, \quad (1)$$

where G is *Newton's gravitational constant*, and c is the speed of light, given by

$$G = 6.67 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2} \quad (2)$$

$$c = 3.00 \times 10^8 \text{ m s}^{-1}. \quad (3)$$

If you venture within R_s of the black hole, you can never escape! You will get a chance to derive this below.

- **Uncertainty.** A second useful fact is *Heisenberg's uncertainty principle*, which tells us that the uncertainty in position, multiplied by the uncertainty in momentum, are at least as big as *Planck's constant* h :

$$\Delta x \cdot \Delta p \geq \frac{h}{4\pi}, \quad h = 6.63 \times 10^{-34} \text{ J s}^{-1}. \quad (4)$$

For classical measurements, this is typically much smaller than the precision of our instruments, but this tiny, inescapable uncertainty has very deep consequences.

- **Relativistic energy.** Third, we need a generalisation of Einstein's famous relation $E = mc^2$, which encompasses both light and matter:

$$E^2 = m^2c^4 + p^2c^2. \quad (5)$$

Here, m is the *rest mass*, and p is the usual classical momentum. This reduces to $E = mc^2$ for a stationary object, but gives

$$E = pc \quad (6)$$

for an object like a photon which has no rest mass.

- **Thermal energy.** Fourth on our list is the connection between temperature and typical energy. For a lump of particles (including massless particles like photons!) at temperature T , the *typical energy per particle* is

$$E_{\text{avg}} \sim k_B T, \quad k_B = 1.38 \times 10^{-23} \text{ J K}^{-1}, \quad (7)$$


where k_B is *Boltzmann's constant*, not to be confused with the (related) Stefan-Boltzmann constant governing luminosity. There are various dimensionless constants that can appear in (7), related to the properties of the system, but we won't need them here.

- **Luminosity.** Finally, we require the *Stefan-Boltzmann law*. This tells us that a hot object at temperature T (in Kelvin), with surface area A , loses energy at a rate L (for *luminosity*) given by

$$L = \sigma AT^4, \quad \sigma = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}, \quad (8)$$

where σ is the *Stefan-Boltzmann constant*.¹

In 1974, Stephen Hawking discovered a remarkable fact: black holes glow, emitting faint radiation like a lump of coal or a light bulb at some temperature T . We can determine the temperature T , and hence the rough time scale for evaporation, but first we should explain how it is even *possible* for a black hole to glow, when by definition it traps light.



The heuristic explanation is as follows: quantum mechanics allows for the production of *virtual pairs of photons* moving in opposite directions. Usually, these pop into existence briefly and then disappear again. But just outside the black hole, one of these photons can fall into the event horizon, while the other zooms off to infinity! It is this second photon that we can detect.

1. Suppose a small particle of mass m is a distance r from the black hole, mass M . The *gravitational potential energy* is

$$U = -\frac{GMm}{r}.$$

This is negative, since we must put energy U into the system to separate the mass m and the black hole so that they no longer feel each other's gravitational influence.

- (a) Suppose we try to separate the particle from the black hole by giving it some kinetic energy. Show that, starting at distance r , it must travel at speed

$$v_{\text{esc}} = \sqrt{\frac{2GM}{r}}$$

in order to escape the black hole's gravitational pull.

- (b) As we get closer to the black hole, the escape velocity will increase. Show that at the Schwarzschild radius (1), the escape velocity becomes the speed of light. Not even photons can escape!
2. (a) For particles produced near the horizon of a black hole, what is the rough uncertainty in position? Don't worry about numerical factors.
 - (b) Using the Heisenberg uncertainty principle (4), argue that the uncertainty in momentum is order

$$\Delta p \sim \frac{hc^2}{GM}.$$

¹Depending on properties of the body and its surrounding medium, the rate L can be modified by an *emissivity factor* ϵ , $0 < \epsilon \leq 1$. A *perfect blackbody* (perfect absorber and emitter) has $\epsilon = 1$. A black hole is, classically, a perfect absorber, and once we take quantum mechanics into account, a perfect emitter too!

3. (a) From Einstein's relation (6), find the uncertainty in energy ΔE . Assuming that the uncertainty in energy is roughly the same as the energy of a typical virtual photon,²

$$E_{\text{avg}} \sim \Delta E,$$

conclude that

$$E_{\text{avg}} \sim \frac{hc^3}{GM}.$$

- (b) From (7), deduce that the Hawking temperature of a black hole of mass M is

$$T_H \sim \frac{hc^3}{k_B GM}. \quad (9)$$

Notice that the black hole gets hotter as it gets smaller!

4. (a) What is the total energy content of a black hole, of mass M , at rest? *Hint.* Use (5).
 (b) The luminosity is the rate of energy loss. Argue (from dimensional analysis or otherwise) that the lifetime of a black hole is

$$t_{\text{life}} \sim \frac{Mc^2}{L} = \frac{Mc^2}{\sigma AT_H^4}.$$

- (c) Use $A \sim R_s^2$ and (9) to show that

$$t_{\text{life}} \sim \frac{G^2 k_B^4 M^3}{\sigma h^4 c^6}. \quad (10)$$

- (d) The sun has mass

$$M_\odot = 2 \times 10^{30} \text{ kg}.$$

Plug this into (10), along with values for the various fundamental constants, and estimate the lifetime of a solar-mass black hole. Give your answer in years, and compare to the age of the universe, $\sim 10^{10}$ y.

²This is a weird property. Usually, the average energy E_{avg} is completely unrelated to the uncertainty ΔE . It just so happens that, for a hot gas of photons, this relation $E_{\text{avg}} \sim \Delta E$ is true. So our assumption is really that the virtual photons form a gas.